arXiv:1609.02437v1 [math.DG] 8 Sep 2016

SURFACES OF CONSTANT CURVATURE IN A SEMI-ISOTROPIC SPACE

MUHITTIN EVREN AYDIN

ABSTRACT. In this paper, we introduce a Lorentz-Minkowski version of the isotropic space which we call it semi-isotropic space. We establish the local theory of non-null curves and surfaces immersed in the semi-isotropic space and provide the general setting in order to study such objects with constant curvature. We describe the surfaces of revolution with constant semi-isotropic mean (H) and semi-relative curvature (K) in the semi-isotropic space. As a further study, we find the surfaces of revolution that satisfy the relation $H^2 = K$.

1. INTRODUCTION

Let $P\left(\mathbb{R}^3\right)$ denote the real projective 3-space. The Cayley-Klein 3-spaces can be defined in $P\left(\mathbb{R}^3\right)$, for example; (semi-) Euclidean, (pseudo-) Galilean, (quasi-) elliptic, (quasi-) hyperbolic and (double-) isotropic space (see [13]).

In particular, the *isotropic 3-space* $\mathbb{I}^3$ has the absolute figure (ω, f_1, f_2), where ω is a plane in $P\left(\mathbb{R}^3\right)$ and f_1, f_2 pair of complex conjugate lines in ω. The *group of motions* of $\mathbb{I}^3$ is a six-parameter group given in the affine coordinates by

$$(\mathbf{x},\mathbf{y},\mathbf{z}) \longmapsto (\mathbf{x}',\mathbf{y}',\mathbf{z}') : \begin{cases} \mathbf{x}' = a + \mathbf{x}\cos\phi - \mathbf{y}\sin\phi, \\ \mathbf{y}' = b + \mathbf{x}\sin\phi + \mathbf{y}\cos\phi, \\ \mathbf{z}' = c + d\mathbf{x} + e\mathbf{y} + \mathbf{z}, \end{cases}$$

where $a, b, c, d, e, \phi \in \mathbb{R}$ (see [24, 29, 30]).

The *isotropic distance* d_i of two points $p = (p_1, p_2, p_3)$ and $q = (q_1, q_2, q_3)$ in $\mathbb{I}^3$ is defined as

$$d_i(p,q) = \sqrt{(q_1 - p_1)^2 + (q_2 - p_2)^2}.$$

The *isotropic scalar product* in $\mathbb{I}^3$ is given by

$$\langle u, v\rangle = \begin{cases} u_1v_1 + u_2v_2, & u_i \neq 0 \text{ or } v_i \neq 0\ ,\ i = 1,2, \\ u_3v_3, & u_i = 0 \text{ and } v_i = 0,\ i = 1,2 \end{cases}$$

for $u = (u_1, u_2, u_3)$ and $v = (v_1, v_2, v_3) \in \mathbb{I}^3$. For more details of $\mathbb{I}^3$ we refer to [1, 2, 6, 12, 16, 23, 27].

The differential geometry of the isotropic spaces was introduced by D. Palman [28], H. Sachs [31] and K. Strubecker [32]. We found motivation for this paper in B. Divjak's works ([9, 10]), where the author introduced the differential geometry of curves and surfaces in the pseudo-Galilean space as generalizing that of the Galilean

2000 *Mathematics Subject Classification.* 53A35, 53B25, 53B30, 53C42.

Key words and phrases. Semi-isotropic space, semi-relative curvature, semi-isotropic mean curvature, surface of revolution, Euler inequality.

space. Intending a similar approach for the isotropic geometry, we establish the Lorentz-Minkowski version of the isotropic space and call it *semi-isotropic space*.

The isotropic space, so called simple isotropic space, is denoted by the notations $\mathbb{I}^3$, I_3^1 or $I_3^{(1)}$ in literature. To avoid confusion, we use the notation $\mathbb{SI}^3$ to specify the semi-isotropic space.

More precisely, the semi-isotropic space $\mathbb{SI}^3$ is indeed the product of the Lorentz-Minkowsi plane $\mathbb{R}_1^2$ and the isotropic line equipped with a degenerate parabolic distance metric, where $\mathbb{R}_1^2$ denotes the Cartesian plane endowed with the metric given by

$$\langle u, v\rangle = u_1 v_1 - u_2 v_2,\ u = (u_1, u_2)\,,\ v = (v_1, v_2) \in \mathbb{R}_1^2.$$

We call it *Lorentzian metric* (also called *Minkowski metric*) and this is a non-degenerate metric of index 1 (see [18, 19, 26]).

The notions such as the norm, the isometric transformations, the causal character of the (non-isotropic) vectors, etc. in the sense of semi-isotropic space are given. We define the planes, the circles and the spheres. Morever, we provide the local theory of non-null curves and surfaces immersed in semi-isotropic space.

On the other hand, an interesting problem in classical differential geometry is to classify surfaces of constant mean (H) and Gaussian curvature (K), obtained by a motion of curves around an axes (no intersect the curve). These are called *surfaces of revolution*. This problem has been solved in various ambient spaces, see [3, 4, 8, 11, 14, 15, 17, 22, 25].

We obtain the surfaces of revolution in $\mathbb{SI}^3$ with constant semi-relative and semi-isotropic mean curvature in this paper.

Another interesting problem is to find the surfaces that satisfy the relation $H^2 = K$. This relation is the equality case of famous Euler inequality for the surfaces in the Euclidean 3-space $\mathbb{R}^3$, i.e.,

$$H^2 \geq K.$$

A surface in $\mathbb{R}^3$ holds $H^2 = K$ if and only if it is totally umbilical, i.e. a part of a plane or a 2-sphere in $\mathbb{R}^3$. For more generalizations of this inequality, see [5, 20, 21]. We also classify the surfaces of revolution in $\mathbb{SI}^3$ with $H^2 = K$.

2. Semi-isotropic space

The *semi-isotropic space* $\mathbb{SI}^3$ is an affine 3-space $\mathbb{R}^3$ endowed with the (semi-) norm defined as

$$\|u\| := \begin{cases} \sqrt{\left|(u_1)^2 - (u_2)^2\right|}, & \text{if } u_1 \neq 0 \text{ or } u_2 \neq 0, \\ |u_3|\,, & \text{if } u_1 = u_2 = 0,\ u = (u_1, u_2, u_3) \in \mathbb{SI}^3. \end{cases}$$

The group of motions of $\mathbb{SI}^3$ is based a six-parameter group G_6 of affine transformations $(\mathbf{x}, \mathbf{y}, \mathbf{z}) \longmapsto (\mathbf{x}', \mathbf{y}', \mathbf{z}')\,,$

$$\begin{cases} \mathbf{x}' := a + \mathbf{x}\cosh\phi + \mathbf{y}\sinh\phi \\ \mathbf{y}' := b + \mathbf{x}\sinh\phi + \mathbf{y}\cosh\phi \\ \mathbf{z}' := c + d\mathbf{x} + e\mathbf{y} + \mathbf{z} \end{cases} \tag{2.1}$$

where $a, b, c, d, e, \phi \in \mathbb{R}$. We call such transformations *semi-isotropic congruence transformations* or *(s-i)-motions.*

Note that (s-i)-motions are the composition an affine shear transformation in **z**-direction and a Lorentzian motion in **xy**−plane. By a Lorentzian motion, we mean a translation and the following Lorentzian rotation

$$\begin{pmatrix} \cosh\phi & \sinh\phi \\ \sinh\phi & \cosh\phi \end{pmatrix},$$

which is one of the four kind isometries of the Lorentz-Minkowski plane $\mathbb{R}^2_1$. In the sequel, many of metric properties in semi-isotropic geometry (invariants under (2.1)) are Lorentzian invariants in the their projections onto $\mathbb{R}^2_1$.

The *semi-isotropic scalar product* between two vectors $u = (u_1, u_2, u_3)$ and $v = (v_1, v_2, v_3) \in \mathbb{SI}^3$ is given by

$$\langle u, v\rangle := \begin{cases} u_1v_1 - u_2v_2, & \text{if at least one of } u_i \text{ or } v_i \text{ is nonzero, } i = 1, 2, \\ u_3v_3, & \text{if } u_i = v_i = 0, \ i = 1, 2. \end{cases}$$

The *vector product* in the sense of semi-isotropic space is

$$u \times v := \begin{vmatrix} e_1 & -e_2 & 0 \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

for $e_1 = (1, 0, 0)$ and $e_2 = (0, 1, 0)$. It can be easily check that

$$\langle u \times v, w\rangle := \det(u, v, \tilde{w}),$$

where $\tilde{w}$ denotes the canonical projection of w onto $\mathbb{R}^2_1$.

We call the vectors of the form $u = (0, 0, u_3)$ in $\mathbb{SI}^3$ *isotropic vectors* and ones of the form $u = (\tilde{u} \neq 0, u_3)$ in $\mathbb{SI}^3$ *non-isotropic vectors*. A vector $u \in \mathbb{SI}^3$ is respectively called *spacelike*, *timelike* and *null* (or *lightlike*) if $\langle u, u\rangle > 0$ or $u = 0$, $\langle u, u\rangle < 0$ and $\langle u, u\rangle = 0$ $(u \neq 0)$.

We remark that only non-isotropic vectors have causal character which is the property to be spacelike, timelike or null.

The set of all null vectors of $\mathbb{SI}^3$ is called *null-cone*, i.e.,

$$\mathcal{C} := \left\{ (x, y, z) \in \mathbb{SI}^3 \middle| \, x^2 - y^2 = 0 \right\} - \left\{ 0 \in \mathbb{SI}^3 \right\}.$$

Timelike-cone is the set of all timelike vectors of $\mathbb{SI}^3$,

$$\mathcal{T} := \left\{ (x, y, z) \in \mathbb{SI}^3 \middle| \, x^2 - y^2 < 0 \right\}.$$

The *semi-isotropic angle* of two timelike non-isotropic vectors $u, v \in \mathbb{SI}^3$ lying in the same timelike-cone is defined as the Lorentzian angle between their projections onto $\mathbb{R}^2_1$, i.e.,

$$\langle u, v\rangle := -\|u\| \, \|v\| \cosh\phi.$$

For a spacelike plane Γ determined by the non-isotropic vectors u, v the induced metric on Γ is positive definite and hence the angle between u and v is the usual Euclidean angle between $\tilde{u}$ and $\tilde{v}$.

Note that all isotropic vectors are orthonogal to non-isotropic ones. Further, two non-isotropic vectors u, v in $\mathbb{SI}^3$ are orthonogal if $\langle u, v\rangle = 0$.

3. Spacelike and timelike curves in $\mathbb{SI}^3$

Let $\alpha(s) = (x(s), y(s), z(s))$ be a regular curve in $\mathbb{SI}^3$, i.e. $\alpha'(s) := \frac{d\alpha}{ds} \neq 0$ for all s. Then it is said to be *admissible* if $\alpha(s)$ has no isotropic tangent vector, i.e. $\tilde{\alpha}'(s) = (x'(s), y'(s)) \neq 0$. An admissible curve $\alpha(s)$ in $\mathbb{SI}^3$ is said to be *spacelike* (resp. *timelike*, *null*) if $\alpha'(s)$ is spacelike (resp. timelike, null) for all s.

We consider only admissible curves whose tangent and normal vector fields are non-null.

Now let $\alpha = \alpha(s)$ be a spacelike curve in $\mathbb{SI}^3$ parameterized by arc-length. Then we have

$$\langle \alpha', \alpha' \rangle = (x')^2 - (y')^2 = 1 \tag{3.1}$$

and taking derivative of (3.1) gives

$$x'x'' - y'y'' = 0. \tag{3.2}$$

Denote $T := \alpha'$ and call it *tangent vector field.* Since $T' = \alpha''$ is timelike in $\mathbb{SI}^3$ we can define the following

$$\kappa := \sqrt{(y'')^2 - (x'')^2},$$

called *semi-isotropic curvature* of α. Using (3.2), we get

$$\kappa := \frac{y''}{x'} \text{ or } \kappa := \frac{x''}{y'}. \tag{3.3}$$

Considering (3.1) and (3.2) into (3.3) we find

$$\kappa := \det\left(\tilde{\alpha}', \tilde{\alpha}''\right). \tag{3.4}$$

Define the *normal vector field* and the *semi-isotropic torsion* of α respectively as

$$N := \frac{1}{\kappa} T' \text{ and } \tau := \frac{\det(\alpha', \alpha'', \alpha''')}{\kappa^2}. \tag{3.5}$$

Since $B := (0, 0, 1)$ is perpendicular to T and N, we can take it as the *binormal vector field of* α.

From (3.5) we have

$$N' = \left(\frac{1}{\kappa}\right)' (x'', y'', z'') + \frac{1}{\kappa} (x''', y''', z'''). \tag{3.6}$$

Put $N' = (n_1, n_2, n_3)$. Hence we write

$$n_1 = \left(\frac{1}{\kappa}\right)' x'' + \frac{1}{\kappa} x'''. \tag{3.7}$$

Using (3.4) into (3.7) yields

$$n_1 = -\frac{x'}{\kappa^2} (x''y''' - x'''y''). \tag{3.8}$$

By taking derivative of (3.2) and considering into (3.8) we obtain

$$n_1 = -\kappa x'. \tag{3.9}$$

Similar computations gives

$$n_2 = -\kappa y'. \tag{3.10}$$

For the third component of N', we have

$$n_3 = \left(\frac{1}{\kappa}\right)' z'' + \frac{1}{\kappa} z'''.$$

It follows from (3.4) that

$$n_3 = \frac{1}{\kappa^2}\left\{-\left(x'y''' - x'''y'\right) z'' + \left(x'y'' - x''y'\right) z'''\right\}. \tag{3.11}$$

By adding and substracting $\left(x''y''' - y''x'''\right) z'$ in (3.11) we conclude

$$n_3 = \frac{1}{\kappa^2}\left\{\det\left(\alpha', \alpha'', \alpha'''\right) - \left(x''y''' - x'''y''\right) z'\right\}. \tag{3.12}$$

Taking derivative of (3.2) and considering into (3.12) implies

$$n_3 = \tau - \kappa z'. \tag{3.13}$$

(3.9) , (3.10) and (3.13) yield that $N' = -\kappa T + \tau B$. Thus we obtain the formulas analogous to these of Frenet as follows

$$\begin{cases} T' := \kappa N \\ N' := -\kappa T + \tau B \\ B' := 0. \end{cases}$$

With similar above arguments we can find the derivative formulas of the vector fields T, N, B for a timelike curve in $\mathbb{SI}^3$ as

$$\begin{cases} T' := \kappa N \\ N' := \kappa T - \tau B \\ B' := 0, \end{cases}$$

where $\kappa := \sqrt{\left(x''\right)^2 - \left(y''\right)^2}$ and $\tau := \frac{\det\left(\alpha', \alpha'', \alpha'''\right)}{\kappa^2}$.

Example 3.1. Consider a hyperbolic cylindrical curve in $\mathbb{SI}^3$ given by (see [7])

$$\alpha(s) = \left(\cosh s, \sinh s, z(s)\right). \tag{3.14}$$

This is a timelike curve of arc-length in $\mathbb{SI}^3$ with $\kappa(s) = 1$ and

$$\tau(s) = z'(s) - z'''(s). \tag{3.15}$$

If we assume that α has constant torsion τ_0, then solving the ordinary differential equation (ODE) (3.15) we find

$$z(s) = \tau_0 s + c_1 e^s - c_2 e^{-s} + c_3,$$

(see Figure 1). This gives the following result.

Proposition 3.1. *Let α be a hyperbolic cylindrical curve in $\mathbb{SI}^3$ with constant torsion τ_0. Then it is of the form*

$$\alpha(s) = \left(\cosh s, \sinh s, \tau_0 s + c_1 e^s - c_2 e^{-s} + c_3\right).$$



FIGURE 1. A hyperbolic cylindrical curve with $\tau_0 = 1$.

4. Planes, circles and spheres in $\mathbb{SI}^3$

The lines in $\mathbf{z}-$direction are called *isotropic lines*. The planes containing an isotropic line are called *isotropic planes*. Other planes are *non-isotropic*.

In the non-isotropic planes the Lorentzian metric is basically used. Let Γ be a non-isotropic plane and $c_0 = (x_0, y_0, z_0)$ a fixed point in Γ. The iso-distance set of c_0 in Γ is

$$\left\{p = (x, y, z) \in \mathbb{SI}^3 : \|p - c_0\| = r, r \in \mathbb{R}^+\right\}.$$

The projection of such a set onto $\mathbf{xy}-$plane is a rectangular hyperbola. We call it a *semi-isotropic circle* (or *(s-i)-circle) of hyperbolic type*.

In the isotropic planes we have an isotropic metric. The *(s-i)-circle of parabolic type* is a parabola with $\mathbf{z}-$axis lying in an isotropic plane. An (s-i)-circle of parabolic type is not the iso-distance set of a fixed point.

We have two types of semi-isotropic spheres. One is the *semi-isotropic sphere* (or *(s-i)-sphere*) *of cylindrical type* which is the set of all points $p \in \mathbb{SI}^3$ with $\|p - c_0\| = r$. This sphere is a right Lorentz hyperbolic cylinder with $\mathbf{z}-$parallel rulings from the Lorentz-Minkowski perspective (see Figure 2).



FIGURE 2. A (s-i)-sphere of cylindrical type.

Other one is the *(s-i)-sphere of parabolic type,*

$$z := r\left(x^2 - y^2\right) + ax + by + c,\ r > 0.$$

These are indeed hyperbolic paraboloids, see Figure 3.

The intersections of these (s-i)-spheres with non-isotropic (resp. isotropic) planes are (s-i)-circles of hyperbolic type (resp. of parabolic type).



FIGURE 3. A (s-i)-sphere of parabolic type.

5. Spacelike and timelike surfaces in $\mathbb{SI}^3$

Let M be a surface immersed in $\mathbb{SI}^3$ without isotropic tangent planes. Then we call such a surface *admissible*. Denote g the metric on M induced from $\mathbb{SI}^3$. The surface M is said to be *spacelike* (resp. *timelike*, *null*) if g is positive definite (resp. a metric with index 1, degenerate).

Throughout this paper we consider only spacelike and timelike admissible surfaces in $\mathbb{SI}^3$.

Assume that M has a local parameterization in $\mathbb{SI}^3$ as follows

$$r : D \subseteq \mathbb{R}^2 \longrightarrow \mathbb{SI}^3 : \ (u_1, u_2) \longmapsto (x(u_1, u_2), y(u_1, u_2), z(u_1, u_2)).$$

If (g_{ij}) is the matrical expression of g with respect to the basis $\{r_{u_1}, r_{u_2}\}$, then we have

$$g_{ij} := \langle r_{u_i}, r_{u_j} \rangle, \ r_{u_i} = \frac{\partial r}{\partial u_i}, \ i, j = 1, 2.$$

The metric g is positive definite if and only if $\det(g_{ij}) > 0$. If the surface M is timelike then $\det(g_{ij}) < 0$.

If M is a graph surface in $\mathbb{SI}^3$ of the form

$$r(u_1, u_2) = (u_1, u_2, z(u_1, u_2)) \tag{5.1}$$

then the metric on M induced from $\mathbb{SI}^3$ is $g = du_1^2 - du_2^2$ and it always becomes a flat timelike surface. So, its Laplacian turns to

$$\triangle := \frac{\partial^2}{\partial u_1^2} - \frac{\partial^2}{\partial u_2^2}.$$

The unit normal vector field of M is the isotropic vector $(0, 0, 1)$ since it is perpendicular to all non-isotropic vectors.

The components of the second fundamental form are

$$h_{ij} := \frac{\det(r_{u_i u_j}, r_{u_1}, r_{u_2})}{\sqrt{|\det(g_{ij})|}}, \ r_{u_i u_j} = \frac{\partial^2 r}{\partial u_i \partial u_j}, \ i, j = 1, 2.$$

For the surfaces of the form (5.1), these components are

$$h_{ij} := z_{u_i u_j} = \frac{\partial^2 z}{\partial u_i \partial u_j}, \ i, j = 1, 2.$$

Thus the *semi-relative curvature* and the *semi-isotropic mean curvature* of M are respectively defined by

$$K := -\varepsilon \frac{\det(h_{ij})}{\det(g_{ij})}$$

and

$$H := -\varepsilon \frac{g_{11}h_{22} - 2g_{12}h_{12} + g_{22}h_{11}}{2\det(g_{ij})},$$

where $\varepsilon := sgn(\det(g_{ij}))$. If M is a graph surface of the form (5.1) then these formulas reduce to

$$K := -\det\left(z_{u_iu_j}\right) \text{ and } H := \frac{1}{2}\triangle z = \frac{z_{u_1u_1} - z_{u_2u_2}}{2}. \tag{5.2}$$

For convenience we abbreviate names of these curvatures as (s-r)-curvature and (s-im)-curvature, respectively.

We call a surface *semi-isotropic flat* or *(s-i)-flat* (resp. *semi-isotropic minimal* or *(s-i)-minimal*) in $\mathbb{SI}^3$ if K (resp. H) vanishes.

According to (5.2) a (s-i)-flat graph surface in $\mathbb{SI}^3$ corresponds to the graph of a solution of homogeneous Monge-Ampère equation, i.e., $\det\left(z_{u_iu_j}\right) = 0$. In case (s-i)-minimal ones these become the graphs in $\mathbb{SI}^3$ associated harmonic functions $\triangle z = 0$.

6. Constant curvature surfaces of revolution in $\mathbb{SI}^3$

Reconsider the (s-i)-motions given by (2.1). Then the Lorentzian rotations in $\mathbb{SI}^3$ are given by, in affine coordinates,

$$\begin{cases} \mathbf{x}' := \mathbf{x}\cosh\phi + \mathbf{y}\sinh\phi \\ \mathbf{y}' := \mathbf{x}\sinh\phi + \mathbf{y}\cosh\phi \\ \mathbf{z}' := \mathbf{z}, \quad \phi \in \mathbb{R}. \end{cases} \tag{6.1}$$

Let $u \longmapsto (u, 0, f(u))$ be a spacelike admissible curve lying in the isotropic $\mathbf{xz}-$plane of $\mathbb{SI}^3$ for a smooth function f. Rotating it around $\mathbf{z}-$axis via Lorentzian rotations given by (6.1) we derive

$$r(u,v) := (u\cosh v, u\sinh v, f(u)). \tag{6.2}$$

We call the rotating curve *profile curve*.

If the profile curve is a timelike curve $u \longmapsto (0, u, f(u))$ lying in the isotropic $\mathbf{yz}-$plane of $\mathbb{SI}^3$, then rotating it around $\mathbf{z}-$axis yields

$$r(u,v) := (u\sinh v, u\cosh v, f(u)). \tag{6.3}$$

The surfaces given by (6.2) and (6.3) are called *surfaces of revolution* in $\mathbb{SI}^3$. Such surfaces of $\mathbb{SI}^3$ are timelike since $\det(g_{ij}) = -u^2$.

The (s-r)-curvature K of these surfaces in $\mathbb{SI}^3$ is

$$K = -\frac{f'f''}{u}, \tag{6.4}$$

where $f'(u) = \frac{df}{du}$, etc.

Now we assume that the surface of revolution has constant (s-r)-curvature K_0 in $\mathbb{SI}^3$. Then (3.4) can be rewritten as

$$f' = \sqrt{c_1 - K_0u^2}, c_1 \in \mathbb{R} \tag{6.5}$$

After integrating (6.5), we obtain

$$f(u) = \frac{c_1}{2\sqrt{K_0}}\arctan\left(\frac{\sqrt{K_0}u}{\sqrt{c_1 - K_0u^2}}\right) + \frac{1}{2}u\sqrt{c_1 - K_0u^2} + c_2, c_2 \in \mathbb{R}$$

which implies the following results.

Theorem 6.1. *Let M a surface of revolution in $\mathbb{SI}^3$ with constant (s-r)-curvature K_0. Then its profile curve is of the form $(u, 0, f(u))$ or $(0, u, f(u))$, where*

$$f(u) = \frac{c_1}{2\sqrt{K_0}} \arctan\left(\frac{\sqrt{K_0}u}{\psi(u)}\right) + \frac{1}{2}u\psi(u) + c_2$$

for $\psi(u) = \sqrt{c_1 - K_0 u^2}$, $c_1, c_2 \in \mathbb{R}$.

We immediately have the following from the Theorem 6.1.

Corollary 6.1. *A surface of revolution is (s-i)-flat in $\mathbb{SI}^3$ if and only if its profile curve is a non-isotropic line which is of the form*

$$(u, 0, c_1 u + c_2) \text{ or } (0, u, c_1 u + c_2),$$

where $c_1, c_2 \in \mathbb{R}$.

The (s-im)-curvature H of a surface of revolution M in $\mathbb{SI}^3$ is

$$H = -\frac{1}{2}\left(\frac{f'}{u} + f''\right). \tag{6.6}$$

Assume that M has constant (s-im)-curvature H_0. After solving the ODE (6.6) we deduce

$$f(u) = -\frac{H_0}{2}u^2 + c_1 \ln u + c_2, \ c_1, c_2 \in \mathbb{R}.$$

Therefore we have proved the following results.

Theorem 6.2. *Let M a surface of revolution in $\mathbb{SI}^3$ with constant (s-im)-curvature H_0. Then its profile curve is of the form $(u, 0, f(u))$ or $(0, u, f(u))$, where*

$$f(u) = -\frac{H_0}{2}u^2 + c_1 \ln u + c_2, \ c_1, c_2 \in \mathbb{R}.$$

Corollary 6.2. *A surface of revolution is (L-I)-minimal in $\mathbb{SI}^3$ if and only if its profile curve is a non-isotropic curve which is of the form*

$$(u, 0, c_1 \ln u + c_2) \text{ or } (0, u, c_1 \ln u + c_2),$$

where $c_1, c_2 \in \mathbb{R}$.

Example 6.1. Take the surfaces of revolution in $\mathbb{SI}^3$ parameterized

$$r(u, v) = (u\cosh v, u\sinh v, u), \ \ (u, v) \in [1, 2] \times [0, 1]$$

and

$$(u, v) = \left(u\cosh v, u\sinh v, \ln u + u^2\right), \ \ (u, v) \in [1, 2] \times [-1, 1].$$

The above first surface is (s-i)-flat and second is a constant (s-im)-curvature surface of revolution, $H = -2$. We plot these as in Figure 4 and Figure 5, respectively.



FIGURE 4. A (s-i)-flat surface of revolution, $K = 0$.



FIGURE 5. A constant (s-i)-mean curvature surface of revolution, $H = -2$.

7. Surfaces of revolution with $H^2 = K$ in $\mathbb{SI}^3$

Next we aim to classify the surfaces of revolution in $\mathbb{SI}^3$ that satisfy $H^2 = K$. For this reason by considering the equalities (6.4) and (6.6) , we have

$$\frac{1}{4}\left(\left(\frac{f'}{u}\right)^2 + 2\frac{f'f''}{u} + (f'')^2\right) = -\frac{f'f''}{u}. \tag{7.1}$$

We can rewirte (7.1) as

$$\left(\frac{f'}{u} + \left(3 - 2\sqrt{2}\right) f''\right)\left(\frac{f'}{u} + \left(3 + 2\sqrt{2}\right) f''\right) = 0,$$

which implies

$$\left(\frac{f'}{u} + \left(3 - 2\sqrt{2}\right) f''\right) = 0 \text{ or } \left(\frac{f'}{u} + \left(3 + 2\sqrt{2}\right) f''\right) = 0.$$

After solving these, we obtain

$$f(u) = -c_1 \frac{u^{-\left(2+2\sqrt{2}\right)}}{2 + 2\sqrt{2}} + c_2 \text{ or } f(u) = c_1 \frac{u^{-2+2\sqrt{2}}}{-2 + 2\sqrt{2}} + c_2$$

for $c_1, c_2 \in \mathbb{R}$.

Theorem 7.1. *Let M a surface of revolution in $\mathbb{SI}^3$ with $H^2 = K$ for (s-im)-curvature H and (s-r)-curvature K. Then its profile curve is of the form $(u, 0, f(u))$ or $(0, u, f(u))$, where*

$$f(u) = -c_1 \frac{u^{-\left(2+2\sqrt{2}\right)}}{2 + 2\sqrt{2}} + c_2 \text{ or } f(u) = c_1 \frac{u^{-2+2\sqrt{2}}}{-2 + 2\sqrt{2}} + c_2, \ c_1, c_2 \in \mathbb{R}. \tag{7.2}$$

Example 7.1. Consider the surface of revolution in $\mathbb{SI}^3$ given via right side equation in (7.2) such that $c_1 = 1$ and $c_2 = 0$. Then its curvatures are of the form

$$K = \left(3 - 2\sqrt{2}\right) u^{-8+4\sqrt{2}},\ H = \left(1 - \sqrt{2}\right) u^{-4+2\sqrt{2}}$$

and satisfy the relation $H^2 = K$. We plot it as in Figure 6.



FIGURE 6. A surface of revolution with $H^2 = K$.

REFERENCES

[1] M.E. Aydin, A Mihai, Ruled surfaces generated by elliptic cylindrical curves in the isotropic space, Georgian Math. J., accepted for publication.

[2] M.E. Aydin, I. Mihai, On certain surfaces in the isotropic 4-space, Math. Commun., in press.

[3] Chr.C. Beneki, G. Kaimakamis, B.J. Papantoniou, A classification of surfaces of revolution with constant Gauss curvature in a 3-dimensional Minkowski space, Bull. Cal. Math. Soc. 90 (1998), 441–458.

[4] R. Caddeo, P. Piu, An. Ratto, Rotational surfaces in 3-dimensional Heisenberg group with constant Gaussian curvature, Boll. Un. Mat. Ital. B(7) 10(2) (1996), 341–357.

[5] B.Y. Chen, Mean curvature and shape operator of isometric immersions in real-space-forms, Glasg. Math. J. 38 (1996), 87–97.

[6] B. Y. Chen, S. Decu, L. Verstraelen, Notes on isotropic geometry of production models, Kragujevac J. Math. 37(2) (2013), 217–220.

[7] M. Crasmareanu, Cylindrical Tzitzeica curves implies forced harmonic oscillators, Balkan J. Geom. Appl., 7(1) (2002), 37-42.

[8] G. Delaunay, Sur la surface de revolution dont la courbure moyenne est constante, J. Math. Pures Appl. Series 6(1) (1841), 309-320.

[9] B. Divjak, Geometrija pseudogalilejevih prostora [Ph.D. thesis], University of Zagreb, 1997.

[10] B. Divjak, Curves in pseudo-Galilean geometry, Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae 41 (1998), 117–128.

[11] M.P. Do Carmo, Differential geometry of curves and surfaces, Prentice Hall: Englewood Cliffs, NJ, 1976.

[12] Z. Erjavec, B. Divjak, D. Horvat, The general solutions of Frenet's system in the equiform geometry of the Galilean, pseudo-Galilean, simple isotropic and double isotropic space, Int. Math. Forum 6(17) (2011), 837-856.

[13] O. Giering, Vorlesungen über höhere Geometrie, Friedr. Vieweg & Sohn, Braunschweig, Germany, 1982.

[14] A. Gray, Modern differential geometry of curves and surfaces with mathematica. CRC Press LLC, 1998.

[15] Z.H. Hou, F. Ji, Helicoidal surfaces with $H^2 = K$ in Minkowski 3-space, J. Math. Anal. Appl. 325 (2007), 101–113.

[16] M.K. Karacan, D.W. Yoon, B. Bukcu, Translation surfaces in the three dimensional simply isotropic space $\mathbb{I}^1_3$, Int. J. Geom. Methods Mod. Phys. 13, 1650088 (2016) (9 pages) DOI: http://dx.doi.org/10.1142/S0219887816500882.

[17] K. Kenmotsu, Surfaces of revolution with prescribed mean curvature, Tohoku Math. J. 32 (1980), 147-153.
[18] R. Lopez, E. Demir, Helicoidal surfaces in Minkowski space with constant mean curvature and constant Gauss curvature, Cent. Eur. J. Math. 12(9) (2014), 1349-1361.
[19] R. Lopez, Differential Geometry of curves and surfaces in Lorentz-Minkowski space, Int. Electron. J. Geom. 7 (2014), 44-107.
[20] A. Mihai, Geometric inequalities for purely real submanifolds in complex space forms, Results Math. 55 (2009), 457–468.
[21] I. Mihai, On the generalized Wintgen inequality for Lagrangian submanifolds in complex space forms, Nonlinear Analysis 95 (2014), 714-720.
[22] Z. Milin-Sipus, B. Divjak, Surfaces of constant curvature in the pseudo-Galilean space, Int. J. Math. Sci., 2012, Art ID375264, 28pp.
[23] Z. Milin-Sipus, B. Divjak, Mappings of ruled surfaces in simply isotropic space I_3^1 that preserve the generators, Monatsh. Math. 139 (2003), 235–245.
[24] Z. Milin-Sipus, Translation surfaces of constant curvatures in a simply isotropic space, Period. Math. Hung. 68 (2014), 160–175
[25] P. Mira, J.A. Pastor, Helicoidal maximal surfaces in Lorentz-Minkowski space, Monatsh. Math. 140 (2003), 315–334.
[26] B. O'Neill, Semi-Riemannian geometry with applications to relativity, Academic Press, New York, 1983.
[27] A.O. Ogrenmis, Rotational surfaces in isotropic spaces satisfying Weingarten conditions, Open Physics 14(9) (2016), 221–225.
[28] D. Palman, Spharische quartiken auf dem torus im einfach isotropen raum, Glasnik Matematicki 14(34) (1979), 345-357.
[29] H. Pottmann, K. Opitz, Curvature analysis and visualization for functions defined on Euclidean spaces or surfaces, Comput. Aided Geom. Design 11 (1994), 655–674.
[30] H. Pottmann, P. Grohs, N.J. Mitra, Laguerre minimal surfaces, isotropic geometry and linear elasticity, Adv. Comput. Math. 31 (2009), 391–419.
[31] H. Sachs, Isotrope Geometrie des Raumes, Vieweg Verlag, Braunschweig, 1990.
[32] K. Strubecker, Differentialgeometrie des isotropen Raumes III, Flachentheorie, Math. Zeitsch. 48 (1942), 369-427.

Department of Mathematics, Faculty of Science, Firat University, Elazig, 23119, Turkey

E-mail address: meaydin@firat.edu.tr